

Scallop (Queen) Dredge on Seagrass (SACs)

Introduction

The Assessing Welsh Fisheries Activities Project is a structured approach to determine the impacts from current and potential fishing activities, from licensed and registered commercial fishing vessels, on the features of Marine Protected Areas.

1. Gear and Feature	Scallop (Queen) Dredge on Seagrass (SACs)
2. Risk Level	Purple (High risk)
3. Description of Feature	<p>Seagrass beds are comprised of several relevant biotopes (see annex 1 for full biotope descriptions).</p> <p>Intertidal seagrass beds biotope LS.LMp.LSgr (and its sub-biotope LS.LMp.LSgr.Znol) are typically dominated by <i>Zostera nolteii</i>.</p> <p>Subtidal seagrass beds biotope SS.SMp.SSgr has sub-biotopes SS.SMp.SSgr.Zmar (dominated by <i>Zostera marina/angustifolia</i> (Note: the taxonomic status of <i>Z. angustifolia</i> is currently under consideration, currently <i>Z. angustifolia</i> is considered a synonym of <i>Z. marina</i>) and SS.SMp.SSgr.Rup (featuring <i>Ruppia maritima</i>).</p> <p>Seagrass beds develop in intertidal and shallow subtidal areas on sands and muds. They may be found in marine inlets and bays but also in other areas, such as lagoons and channels, which are sheltered from significant wave action (BRIG, 2008).</p> <p>The <i>Zostera</i> species that occur in the UK all are considered to be scarce. Dwarf eelgrass <i>Zostera nolteii</i> is found highest on the shore, often adjacent to lower saltmarsh communities. Narrow-leaved eelgrass <i>Zostera marina</i> is found on the mid to lower shore and in the sublittoral. The plants stabilise the substratum, are an important source of organic matter and provide shelter and a surface for</p>

attachment by other species.

Eelgrass is an important source of food for wildfowl which feed on intertidal beds. Where this habitat is well developed the leaves of eelgrass plants may be colonised by diatoms and algae such as *Ulva* spp., *Cladophora* spp., Red Seagrass Crust *Rhodophysema georgii*, *Ceramium virgatum*, stalked jellyfish and anemones. The soft sediment infauna may include amphipods, polychaete worms, bivalves and echinoderms.

The shelter provided by seagrass beds makes them important nursery areas for flatfish and, in some areas for cephalopods. Adult fish frequently seen in *Zostera* beds include pollack *Pollachius pollachius*, two-spotted goby *Gobiusculus flavescens* and various wrasse species (BRIG, 2008; Bertelli & Unsworth, 2014). Two species of pipefish, *Entelurus aequoraeus* and *Syngnathus typhie* are almost totally restricted to seagrass beds while the red algae *Polysiphonia harveyi* which has only recently been recorded from the British Isles is often associated with eelgrass beds (BRIG, 2008).

The diversity of species associated with the seagrass bed will depend on environmental factors such as salinity and tidal exposure and the density of microhabitats, but it is potentially highest in the perennial fully marine subtidal communities and may be lowest in intertidal, estuarine, annual beds (BRIG, 2008).

The diversity of species associated with the seagrass bed will depend on environmental factors such as salinity and tidal exposure and the density of microhabitats, but it is potentially highest in the perennial fully marine subtidal communities and may be lowest in intertidal, estuarine, annual beds (BRIG, 2008).

Zostera beds are naturally dynamic and may show marked seasonal changes. Leaves are shed in winter, although *Zostera noltii* retains its leaves longer than *Zostera marina*. Leaf growth stops in September/October (Brown, 1990).

	<p>Although a wide range of species are associated with seagrass beds which provide habitat and food resources, these species occur in a range of other biotopes and were therefore not considered to characterize the sensitivity of this biotope (D'Avack <i>et al</i>, 2014).</p> <p>Seagrass species are fast-growing and relatively short-lived, they can take a considerable time to recover from damaging events, if recovery does occur at all (D'Avack <i>et al</i>, 2014).</p> <p>Boese <i>et al</i> (2009) found that natural seedling production was not of significance in the recovery of seagrass beds but that recovery was due exclusively to rhizome growth from adjacent perennial beds. All <i>Zostera</i> plants have a similar type of structure and they are restricted to horizontal growth of roots and, hence, unable to grow rhizomes vertically.</p>
<p>4. Description of Gear</p>	<p>Queen scallops (<i>Aequipecten opercularis</i>) are predominantly targeted using towed fishing gear, either in the form of skid dredges (modified Newhaven dredges) or modified otter trawls.</p> <p>Queen scallops are more active swimmers than king scallops and do not recess into the seabed (Brand, 2006). Dredges and otter trawls take advantage of the natural propensity of queen scallops to swim up into the water column when disturbed, rather than relying on extraction of the scallops from the sediment as is the case for Newhaven dredges (Beukers-Stewart & Beukers-Stewart, 2009).</p> <p>A modified Newhaven dredge can be about 1.95m wide, often with a higher front opening. Instead of metal teeth it can have a rubber lip or sometimes the front part of the dredge consists of a metal grid mounted on four rubber rollers, two on each side of the grid. Alternatively, the tooth bar is replaced with a tickler chain. The modified dredge is normally fitted with skis or skids on either side designed to run along the top of the seabed. The dredge has a traditional metal belly bag with a mesh size of 60mm to retain the queen scallops (Humphey, 2009).</p>

	<p>Traditional toothed king scallop dredges are occasionally used to target queen scallops, these dredges are approximately 0.76m wide, with a chain mail belly bag and a 60mm mesh. Each dredge bar usually has 17 metal teeth of around 6cm in length on it (Hinz <i>et al</i>, 2009). The amount of dredges per side of the vessel can vary between 1 and 16 depending on the size and power of the vessel.</p> <p>The choice of skid dredges or otter trawls is largely governed by the nature of the substrate on different fishing grounds, with skid dredges being more effective in rough/coarse sediment areas and trawls in sandy/muddy areas (Vause <i>et al</i>, 2007).</p>
<p>5. Assessment of Impact Pathways:</p> <ol style="list-style-type: none"> 1. Damage to a designated habitat feature (including through direct physical impact, pollution, changes in thermal regime, hydrodynamics, light etc). 2. Damage to a designated habitat feature via removal of, or other detrimental impact on, typical species. 	<p>The three types of queen scallop fishing gear described above are all bottom contacting gear and as such impose a similar effect upon the seabed. The assessment below incorporates all three gears and will be referred to collectively as scallop (queen) dredge gear. Any differences in gear interactions will also be described below.</p> <ol style="list-style-type: none"> 1. Demersal mobile fishing gear reduces habitat complexity by: removing emergent epifauna, smoothing sedimentary bedforms, and removing taxa that produce structure (Auster & Langton, 1999). Ways in which gear affects the seabed can be classified as: scraping and ploughing; sediment resuspension; and physical destruction, removal, or scattering of non-target benthos (Jones, 1992). <p>The Newhaven dredges and trawls employed in this fishery are also known to cause considerable damage and disturbance to benthic communities and associated nursery habitat (Eleftheriou & Robertson, 1992; Jennings <i>et al</i>, 2001; Kaiser <i>et al</i>, 2006). The action of the scallop (queen) dredge gear, like the impacts from king scallop dredges, will impact the seagrass by direct physical contact with it and through removal of the surface layer of the sediment. This contact will damage the seagrass and the seagrass' root system.</p> <p>As a sensitive marine habitat, seagrass meadows are highly susceptible to physical impacts and disturbance of the habitat (Short & Wyllie-Echeverria, 1996). Most seagrass species, including <i>Zostera</i></p>

marina and *Zostera noltii*, grow over sandy to muddy sediments, which are easily penetrated by seagrass roots. The direct ploughing and scraping of the scallop (queen) dredge gear on seagrass could cause mortality from a single pass of a scallop dredge. Scallop (queen) dredge gear have penetration depths of 1-15cm in sand and 1-35cm in mud (Eigaard *et al*, 2016; Paschen *et al*, 2000) and could remove the upper layers of sediment on which the seagrasses are reliant for anchoring and nutrient uptake.

Unsworth and Cullen-Unsworth (2015) investigated the effects of physical disturbance on seagrass meadows in Porthdinllaen, within the Pen Llyn a'r Sarnau Special Areas of Conservation. They conclude that the chains and anchors associated to various types of moorings drag over the seagrass and repeatedly tear the plants, eventually ripping up their roots and rhizomes and reducing the capacity for recovery to occur. The effects of towed demersal gear, such as scallop dredge gear, on seagrasses is likely to be greater than the damage caused by anchoring and moorings.

A depression of the seabed caused by disturbance of the sediment can restrict the expansion of the seagrass bed. The size and shape of impacted areas will have a considerable effect on resilience rates (Creed *et al*, 1999). Larger denuded areas (such as those caused by towed demersal fishing gear) are likely to take longer to recover than smaller scars, for example seagrass beds likely to be more resilient to physical damage resulting from narrow furrows left after anchoring because of large edge-to-area ration and related availability of plants for recolonisation.

Neckles *et al* (2005) investigated the effects of trawling for the blue mussels *Mytilus edulis* on *Zostera marina* beds in Maquoit Bay, USA. Impacted sites ranged from 3.4 to 31.8ha in size and were characterized by the removal of above and belowground plant material from the majority of the bottom. The study found that one year after the last trawl, *Zostera marina* shoot density, shoot height and total biomass averaged respectively to 2-3%, 46-61% and < 1% to that of the reference sites. Substantial differences in *Zostera*

marina biomass persisted between disturbed and reference sites up to 7 years after trawling. Rates of recovery depended on initial fishing intensity but the authors estimated that an average of 10.6 years was required for *Zostera marina* shoot density to match pre-trawling standards.

The effects of dredging for scallops on *Zostera marina* beds were investigated by Fonseca *et al* (1984) in Nova Scotia, USA. Dredging was carried out when *Zostera marina* was in its vegetative stage on hard sand and on soft mud substrata. Damage was assessed by analysing the effects of scallop harvesting on seagrass foliar dry weight and on the number of shoots. Lower levels of dredging (15 dredges) had a different impact depending on substrata, with the hard bottom retaining a significantly greater overall biomass than soft bottom. However, an increase in dredging effort (30 dredges) led to a significant reduction in *Zostera marina* biomass and shoot number on both hard and soft bottoms. Solway Firth is a British example for the detrimental effects of dredging on seagrass habitats. In the area, where harvesting for cockles by hand is a traditional practice, suction dredging was introduced in the 1980s to increase the yield. A study by Perkins (1988) found that where suction dredging occurred, the sediment was smoothed and characterized by a total absence of *Zostera* plants. The study concluded that the fishery was causing widespread damage and could even completely eradicate *Zostera* from affected areas. Due to concerns over the sustainability of this fishing activity, the impacts on cockle and *Zostera* stocks, and the effects on overwintering wildfowl, the fishery was closed to all forms of mechanical harvesting in 1994.

Most seagrass species grow over sandy to muddy sediments, which are easily penetrated by seagrass roots. However, highly mobile, but otherwise suitable, sandy sediments may be bare of seagrass (Hemminga & Duarte, 2000). Processes that cause sand ripples and waves can cause successive burial and erosion, which may cause seagrass mortality, depending on the size and frequency of these events. Sediment disturbance caused by scallop dredging is likely to

cause a greater intensity of burial and erosion in a single pass of the gear than caused by current and wave energy. Below ground rizomes and root structures are dependant on the upper few centimeters of sediment for nutrients. Continued scallop dredging events could reduce the nutrient levels within sediments and make recovery difficult. The depth limit of seagrasses is set by the compensation irradiance for growth, or the irradiance required to provide sufficient carbon gains to compensate for carbon losses.

The light requirement for seagrass growth is typically defined as the percentage of surface irradiance that needs to be received by the plants to grow, which ranges between 4% and 29% (Dennison *et al*, 1993), with an average of about 11% of the irradiance incident just below the water surface (Duarte, 1991). These light requirements are greater than those generally observed for other marine phototrophs, such as macroalgae and microalgae (Duarte, 1995). These extremely high light requirements mean that seagrasses are acutely responsive to environmental changes, especially those that alter water clarity (Orth *et al*, 2006).

Duarte *et al* (2007) sought to test seagrass depth limit models from test data comprising 424 reports of seagrass colonisation depth limits. Most (86%) of the reports in the validation set assembled pertained to observations of colonisations depth of *Zostera marina*. The results showed that *Zostera marina* has a colonisation depth range of between 0.5-10m. This data has taken into account varying levels of turbidity. Duarte *et al* (2007) does however make the argument that clear water could allow seagrasses to grow at a depth of 30m. At these depths, the contribution of absorption of water filters out irradiance at red wavelengths while allowing high-energy blue light to penetrate and promote photosynthesis.

Trawling and dredging re-suspend large amounts of sediments (PilskaIn *et al*, 1998). The increase in turbidity through sediment re-suspension caused by scallop (queen) dredging would influence the photosynthesis of seagrasses, which could cause mortality. Riemann and Hoffmann (1991) found short-term increased suspended sediment loads of 960-1361%.

In conclusion, the direct physical impact, changes in light caused by sediment re-suspension or sediment removal by, fishing with scallop (queen) dredge gear could cause seagrass mortality, without guarantee of recolonisation or recovery.

2. Demersal trawls cause direct mortality to non-target organisms through impact on the seabed (Bergman & van Santbrink, 2000).

There is growing evidence that seagrass meadows are presently experiencing worldwide decline primarily because of human disturbance, such as direct physical damage and deterioration of water quality (Short & Wyllie-Echeverria, 1996; Hemminga & Duarte, 2000). There is, therefore, concern that the functions seagrasses perform in the marine ecosystem will be reduced or, in some places, lost altogether (Duarte, 2002). Fisheries operations, particularly shallow trawling (Pascualini *et al*, 1999), causes disturbance and damage to seagrass communities.

Seagrass meadows can serve as a nursery ground, often to juvenile stages of economically important species of finfish and shellfish, although the species vary by region and climate (Beck *et al*, 2001; Heck *et al*, 2003). The loss of seagrasses, through physical disturbance from scallop (queen) dredge gear, would therefore impact on the typical species in which it supports.

Collie *et al* (2000) undertook an analysis of published research into fishing activity impacts on the seabed, based on 39 research projects undertaken previously. They found an average of 46% decrease in total number of individuals of a species in study sites that were disturbed with bottom towed gear.

In conclusion, seagrass loss through scallop (queen) dredge gear could cause a detrimental impact on typical species through loss of food and removal of nursery areas for juvenile finfish and shellfish species. Fishing with scallop (queen) dredge gear could also directly remove typical species from the feature. Typical species

		recolonisation of this habitat would be dependant on the quality of habitat which remained following a dredging episode. Where there is damage to the habitat, mobile species would be quick to recolonise. If there is the total removal of seagrasses, recolonisation will not occur.
6. MPAs where feature exists	Menai Strait and Conwy Bay SAC	Intertidally between Llanfairfechan and Bangor, at Moel-y-Don opp Felinheli and within Y Foryd.
	Lleyn Peninsular and Sarnau SAC	Intertidal and Subtidal beds at Porth Dinllaen, Llanbedrog, intertidally at Pen y chain, subtidally off Criccieth (within 1Nm).
	Pembrokeshire Marine SAC	Subtidally within North Haven at Skomer, intertidally and subtidally within the Milford Haven at Sandy Haven Bay, intertidally on Dale Flats, subtidally and intertidally between South Hook Point and Milford Docks, Sprinkle Pill, Garron Pill, Cresswell River, Carew River, Coshaston Pill, West Llanion Pill, Pembroke River, Pwllcrochan Flats, Angle Bay, off Ellen's Well and the Lifeboat station.
	Carmarthen Bay and Estuaries SAC	River Towy (between Salmon Point and Ferryside), within the Burry Inlet at Llanridian Sands and Penrhyn Gwyn.
	Severn Estuary SAC	Located between Summerleaze and the M4 Severn Crossing.

7. Conclusion

The information presented above indicates that the action of fishing with scallop (queen) dredge gear directly on seagrass (SACs) is likely to cause lethal damage to the seagrass and associated species, while recovery is possible (up to 10.6 years) this would be less likely if the upper centimeters of sediment were removed during the initial interaction or prolonged fishing. Additionally, fishing with scallop (queen) dredge gear adjacent to seagrass beds could have a negative impact from short or long term sediment re-suspension causing an increase in turbidity, thus affecting photosynthesis; this impact would depend on the extent and frequency of the activity and the tidal and environmental conditions in the area of the habitat.

8. References

- Auster, P.J. & Langton, R.W. (1996). The effects of fishing on fish habitat. In: Benaka L (ed) Fish habitat essential fish habitat (EFH) and rehabilitation. *Am Fish Soc* 22:150-187
- Beck, M.W., Heck Jr, K.L., Able, K.W., Childers, D.L., Eggleston, D.B., Gillanders, B.M., Halpern, B., Hays, C.G., Hoshino, K., Minello, T.J., Orth, R.J. (2001). The identification, conservation, and management of estuarine and marine nurseries for fish and invertebrates: a better understanding of the habitats that serve as nurseries for marine species and the factors that create site-specific variability in nursery quality will improve conservation and management of these areas. *Bioscience*, 51(8), pp.633-641.
- Bertelli, C.M. & Unsworth, R.K.F. (2014). Protecting the hand that feeds us: Seagrass (*Zostera marina*) serves as commercial juvenile fish habitat. *Marine Pollution Bulletin*, Vol. 83, Issue 2: 425-429
- Beukers-Stewart B.D. & Beukers-Stewart J.S. (2009). Principles for the management of inshore scallop fisheries around the United Kingdom. Report to Natural England, Countryside Council for Wales and Scottish Natural Heritage. University of York. 57pp.
- Boese, B.L., Kaldy, J.E., Clinton, P.J., Eldridge, P.M. & Folger, C.L. (2009). Recolonization of intertidal *Zostera marina* L. (eelgrass) following experimental shoot removal. *Journal of Experimental Marine Biology and Ecology*, **374** (1), 69-77.
- Brand, A.R. (2006). Scallop Ecology: Distributions and Behaviour. In: Shumway S, Parsons GJ (eds) *Scallops: Biology, Ecology and Aquaculture*. Elsevier, Amsterdam, p 1460
- BRIG. (ed. Ant Maddock) (2008). UK Biodiversity Action Plan Priority Habitat Descriptions: Seagrass Beds (available from <http://jncc.defra.gov.uk/page-5706>)
- Brown, R.A. (1990). Strangford Lough. The wildlife of an Irish sea lough. The Institute of Irish Studies, Queens University of Belfast.
- Collie, J.S., Hall, S.J., Kaiser, M.J. & Poiner, I.R. (2000). A quantitative analysis of fishing impacts shelf-sea benthos. *Journal of Animal Ecology*, 69(5), 785–798.
- Creed, J.C., Filho, A. & Gilberto, M. (1999). Disturbance and recovery of the macroflora of a seagrass *Halodule wrightii* (Ascherson) meadow in the Abrolhos Marine National Park, Brazil: an experimental evaluation of anchor damage. *Journal of Experimental Marine Biology and Ecology*, **235** (2), 285-306.
- D’Avack, E.A.S., Tillin, H., Jackson, E.L. & Tyler-Walters, H. (2014). Assessing the sensitivity of seagrass bed biotopes to pressures associated with marine activities. JNCC Report No. 505. *Joint Nature Conservation Committee*, Peterborough. Available from www.marlin.ac.uk/publications.
- Dennison, W.C., Orth, R.J., Moore, K.A., Stevenson, J.C., Carter, V., Kollar, S., Bergstrom, P.W., Batiuk, R.A. (1993). Assessing water quality with submersed aquatic vegetation. *BioScience*, 43, 86 – 94.
- Duarte, C.M. (1991). Seagrass depth limits. *Aquatic Botany*, 40, 363 – 377.
- Duarte, C.M. (1995). Submerged aquatic vegetation in relation to different nutrient regimes. *Ophelia*, 41, 87 – 112.
- Duarte, C.M. (2002). The future of seagrass meadows. *Environmental Conservation*, 29, 192-206.
- Duarte, C.M., Marba, N., Krause-Jensen, D., Sanchez-Camacho, M. (2007). Testing the predictive power of seagrass depth limit models. *Estuaries and Coasts*, 30, 652-656.

- Eigaard, O.R., Bastardie, F., Breen, M., Dinesen, G.E., Hintzen, N.T., Laffargue, P., Mortensen, L.O., Nielsen, J.R., Nilsson, Hans C., O'Neill, F.G., Polet, H., Reid, D.G., Sala, A., Sko"ld, M., Smith, C., Sorensen, T.K., Tully, O., Zengin, M. & Rijnsdorp, A.D. (2016). Estimating seabed pressure from demersal trawls, seines, and dredges based on gear design and dimensions. – ICES Journal of Marine Science, 73: i27–i43.
- Eleftheriou, A. & Robertson, M.R. (1992). The effects of experimental scallop dredging on the fauna and physical environment of a shallow sandy community. *Neth J Sea Res* 30:289–299
- Fonseca, M.S., Thayer, G.W., Chester, A.J. & Foltz, C. (1984). Impact of scallop harvesting on eelgrass (*Zostera marina*) meadows. *North American Journal of Fisheries Management*, 4 (3), 286-293.
- Heck Jr, K.L., Hays, G., Orth, R.J. (2003). Critical evaluation of the nursery role hypothesis for seagrass meadows. *Marine Ecology Progress Series*, 253, 123-136.
- Hemminga, M.A. & Duarte, C.M. (2000). Seagrass ecology. *Cambridge University Press*.
- Hinz, H., Murray, L.G. & Kaiser, M.J. (2009). Efficiency and environmental impacts of three different Queen scallop fishing gears. Fisheries & Conservation report No. 8, Bangor University. pp.23.
- Humphey, M. (2009). Testing Materials used in Queen Scallop dredge Construction. SEAFISH report: SR612
- Jennings, S., Kaiser, M.J., Reynolds, J.D. (2001). Marine fisheries ecology. Blackwell Science, Oxford
- Jones, B. (1992). Environmental impact of trawling on the seabed: A review, *New Zealand Journal of Marine and Freshwater Research*, 26:1, 59-67,
- Kaiser, M.J., Clarke, K.R., Hinz, H., Austen, M.C.V., Somerfield, P.J., Karakassis, I. (2006). Global analysis of response and recovery of benthic biota to fishing. *Mar Ecol Progr Ser* 311:1–14
- Neckles, H.A., Short, F.T., Barker, S. & Kopp, B.S. (2005). Disturbance of eelgrass *Zostera marina* by commercial mussel *Mytilus edulis* harvesting in Maine: dragging impacts and habitat recovery. *Marine Ecology Progress Series*, 285, 57-73.
- Orth, R.J., Carruthers, T.J., Dennison, W.C., Duarte, C.M., Fourqurean, J.W., Heck Jr, K.L., Hughes, A.R., Kendrick, G.A., Kenworthy, W.J., Olyarnik, S., Short, F.T. (2006). A global crisis for seagrass ecosystems. *Bioscience*, 56(12), pp.987-996.
- Paschen, M., Richter, U. & Ko"pnick, W. (2000). Trawl Penetration in the Seabed (TRAPESE). Final report Contract No. 96–006.
- Pascualini, V., Pergent-Martini, C., Pergent, G. (1999). Environmental impact identification along the Corsican coast (Mediterranean sea) using image processing. *Aquatic Botany*, 65, 311-320.
- Perkins, E.J. (1988). The impact of suction dredging upon the population of cockles *Cerastoderma edule* in Auchencairn Bay. *Report to the Nature Conservancy Council, South-west Region, Scotland*, no. NC 232 I).
- Pilskaln, C.H., Churchill, J.H., Mayer, L.M. (1998). Frequency of bottom trawling in the Gulf of Maine and speculations on the geochemical consequences. *Conservation Biology* 12: 1223-1229
- Riemann, B. & Hoffmann, E. (1991). Ecological consequences of dredging and bottom trawling in the Limfjord, Denmark. *Marine Ecology Progress Series* 69:171-178.
- Short, F.T. & Wyllie-Echeverria, S. (1996). Natural and human-induced disturbances of seagrasses. *Environmental Conservation*, 23, 17-27.

- Unsworth, R.K.F. & Cullen-Unsworth, L.C. (2015). Pen Llyn a'r Sarnau Special Area of Conservation (SAC) Porthdinllaen Seagrass Project: A review of current knowledge. Report for Gwynedd Council.
- Vause, B. J., Beukers-Stewart, B.D. & Brand, A.R. (2007). Fluctuations and forecasts on the fishery for queen scallops (*Aequipecten opercularis*) around the isle of man. *Ices Journal of Marine Science* **64**: 1124-1135.

Annex 1

Biotope descriptions (version 15.03) (JNCC - <http://jncc.defra.gov.uk/marine/biotopes/hierarchy.aspx?level=5>)

LS.LMp.LSgr.Znol - *Zostera noltii* beds in littoral muddy sand

Mid and upper shore wave-sheltered muddy fine sand or sandy mud with narrow-leafed eel grass *Zostera noltii* at an abundance of frequent or above. It should be noted that the presence of *Z. noltii* as scattered fronds does not change what is otherwise a muddy sand biotope. Exactly what determines the distribution of *Z. noltii* is not entirely clear. It is often found in small lagoons and pools, remaining permanently submerged, and on sediment shores where the muddiness of the sediment retains water and stops the roots from drying out. An anoxic layer is usually present below 5 cm sediment depth. The infaunal community is characterised by the polychaetes *Scoloplos armiger*, *Pygospio elegans* and *Arenicola marina*, oligochaetes, the spire shell *Hydrobia ulvae*, and the bivalves *Cerastoderma edule* and *Macoma balthica*. The green algae *Enteromorpha* spp. may be present on the sediment surface. The characterising species lists below give an indication both of the epibiota and of the sediment infauna that may be present in intertidal seagrass beds. Note that the correct spelling is *Z. noltei*. Horneum. Stace 1991 (reprinted 2001).

SS.SMp.SSgr.Zmar - *Zostera marina/angustifolia* beds on lower shore or infralittoral clean or muddy sand

Expanses of clean or muddy fine sand and sandy mud in shallow water and on the lower shore (typically to about 5 m depth) can have dense stands of *Zostera marina/angustifolia* [Note: the taxonomic status of *Z. angustifolia* is currently under consideration]. In Zmar the community composition may be dominated by these *Zostera* species and therefore characterised by the associated biota. Other biota present can be closely related to that of areas of sediment not containing *Zostera marina*, for example, *Laminaria saccharina*, *Chorda filum* and infaunal species such as *Ensis* spp. and *Echinocardium cordatum* (e.g. Bamber 1993). From the available data it would appear that a number of sub-biotopes may be found within this biotope dependant on the nature of the substratum and it should be noted that sparse beds of *Zostera marina* may be more readily characterised by their infaunal community. For example, coarse marine sands with seagrass have associated communities similar to MoeVen, SLan or Glap whilst muddy sands may have infaunal populations related to EcorEns, ArelSa and FfabMag. Muddy examples of this biotope may show similarities to SundAasp, PhiVir, Are or AfilMysAnit. At present the data does not permit a detailed description of these sub-biotopes but it is likely that with further study the relationships between these assemblages will be clarified. Furthermore, whilst the *Zostera* biotope may be considered an epibiotic overlay of established sedimentary communities it is likely that the presence of *Zostera* will modify the underlying community to some extent. For example, beds of this biotope in the south-west of Britain may contain conspicuous and distinctive assemblages of Lusitanian fauna such as *Laomedea angulata*, *Hippocampus* spp. and Stauromedusae. In addition, it is known that seagrass beds play an important role in the trophic status of marine and estuarine waters, acting as an important conduit or sink for nutrients and consequently some examples of *Zostera marina* beds have markedly anoxic sediments associated with them.

SS.SMp.SSgr.Rup – *Ruppia maritima* in reduced salinity infralittoral muddy sand

In sheltered brackish muddy sand and mud, beds of *Ruppia maritima* and more rarely *Ruppia spiralis* may occur. These beds may be populated by fish such as *Gasterosteus aculeatus* which is less common on filamentous algal-dominated sediments. Seaweeds such as *Chaetomorpha* spp., *Enteromorpha* spp., *Cladophora* spp., and *Chorda filum* are also often present in addition to occasional fucoids. In some cases the stoneworts *Lamprothamnium papulosum* and *Chara aspera* occur. Infaunal and epifaunal species may include mysid crustacea,

the polychaete *Arenicola marina*, the gastropod *Hydrobia ulvae*, the amphipod *Corophium volutator* and oligochaetes such as *Heterochaeta costata*. In some areas *Zostera marina* may also be interspersed with the *Ruppia* beds.